

<i>STD/I</i>	<i>What to Watch For</i>	<i>How do you get this STD?</i>	<i>What Happens if you don't get Treated?</i>	<i>Will condoms prevent infection?</i>
<i>Chlamydia or NGU</i>	<ul style="list-style-type: none"> Chlamydia affects women and men. In men, chlamydia can cause NGU. Symptoms show up 7 – 28 days after having sex. Most women and some men have no symptoms. <p>Women</p> <ul style="list-style-type: none"> Discharge from the vagina. Bleeding form the vagina between periods. Burning or pain when you urinate (pee). Need to urinate (pee) more often. Pain in the abdomen, sometimes with fever and nausea. <p>Men</p> <ul style="list-style-type: none"> Watery, white drip from the penis. Burning or pain when you urinate (pee). Need to urinate (pee) more often. Swollen or tender testicles. 	Spread during vaginal, anal or oral sex with someone who has chlamydia.	<ul style="list-style-type: none"> You can give chlamydia to your sex partners. Can lead to a more serious infection. Reproductive organs can be damaged. Women and possibly men may no longer be able to have children. A mother with chlamydia can give it to a child during childbirth. 	YES
<i>Genital Warts (HPV)</i>	<ul style="list-style-type: none"> Symptoms show up 1-8 months after contact with HPV, the virus that causes genital warts. (Another type of HPV is linked to cervical cancer in women.) Small, bumpy warts on the sex organs and anus. Itching or burning around the sex organs. After warts go away, the virus sometimes stays in the body. The warts can come back. 	Spread during vaginal, anal and oral sex, and sometimes by touching an area with warts, with someone who has genital warts (HPV).	<ul style="list-style-type: none"> You can give genital warts to your sex partners. Warts may go away on their own, remain unchanged, or grow and spread. A mother with warts can give them to her baby during childbirth. 	NO
<i>Gonorrhea</i>	<ul style="list-style-type: none"> Symptoms show up 2-21 days after having sex. Most women and some men have no symptoms. <p>Women</p> <ul style="list-style-type: none"> Thick yellow or gray discharge from the vagina. Burning or pain when you urinate (pee) or have a bowel movement. Abnormal periods or bleeding between periods. Cramps and pain in the lower belly. <p>Men</p> <ul style="list-style-type: none"> Thick yellow or greenish drip from the penis. Burning or pain when you urinate (pee) or have a bowel movement. Need to urinate (pee) more often. Swollen or tender testicles. 	Spread during anal and oral sex with someone who has gonorrhea.	<ul style="list-style-type: none"> You can give gonorrhea to your sex partners. Can lead to a more serious infection. Reproductive organs can be damaged. Both men and women may no longer be able to have children. Can cause heart trouble, skin disease, arthritis and blindness. A mother with gonorrhea can give it to her baby in the womb or during childbirth. 	YES
<i>Hepatitis B</i>	<ul style="list-style-type: none"> Symptoms show up 1-9 months after contact with the Hepatitis B virus. Many people have no symptoms or mild symptoms. Flu-like feelings that don't go away. Tiredness. Yellow skin. Dark urine, light colored bowel movements. 	<ul style="list-style-type: none"> Spread during vaginal, anal and oral sex with someone who has Hepatitis B. Spread by sharing needles to inject any substance. Spread by contact with infected blood or blood products. 	<ul style="list-style-type: none"> You can give Hepatitis B to your sex partner(s) or someone you share a needle with. Some people cannot be cured. Can cause permanent liver damage or liver cancer. A mother with Hepatitis B can give it to her baby during childbirth. 	YES

<i>STD/I</i>	<i>What to Watch For</i>	<i>How do you get this STD?</i>	<i>What Happens if you don't get Treated?</i>	<i>Will condoms prevent infection?</i>
<i>Herpes</i>	<ul style="list-style-type: none"> Symptoms show up 1-30 days or longer after having sex. Some people have no symptoms. Flu-like feelings. Small, painful blisters on the sex organs or mouth. Itching or burning before the blisters appear. Blisters last 1-3 weeks. Blisters go away, but you still have herpes. Blisters can come back. 	<ul style="list-style-type: none"> Spread during vaginal, anal and oral sex and sometimes touching an affected area with someone who has herpes. 	<ul style="list-style-type: none"> You can give herpes to your sex partners. Herpes cannot be cured. A mother with herpes can give it to her baby during childbirth. 	NO
<i>HIV/AIDS</i>	<ul style="list-style-type: none"> Symptoms show up several months to several years after contact with HIV, the virus that causes AIDS. Can be present for many years with no symptoms. Unexplained weight loss or tiredness. Flu-like symptoms that don't go away. Diarrhea. White spots in the mouth. In women, yeast infections that won't go away. 	<ul style="list-style-type: none"> Spread during vaginal, anal and oral sex with someone who has HIV. Spread by sharing needles to inject any substance. Spread by contact with infected blood or blood products. 	<ul style="list-style-type: none"> You can give HIV to your sex partner(s) or someone you share a needle with. A mother with HIV can give it to her baby in the womb, during childbirth or while breastfeeding. HIV cannot be cured. Can cause illness and death. 	YES
<i>Syphilis</i>	<p>1st Stage:</p> <ul style="list-style-type: none"> Symptoms show up 1-12 weeks after having sex. Painless sores on the mouth or sex organs. Sores last 1-5 weeks. Sores go away, but you still have syphilis. <p>2nd Stage:</p> <ul style="list-style-type: none"> Symptoms show up 1 week – 6 months after sores heal. A rash anywhere on the body. Flu-like feelings. Rash and flu-like feelings go away, but you still have syphilis. 	<p>Spread during vaginal, anal and oral sex with someone who has syphilis.</p>	<ul style="list-style-type: none"> You can give syphilis to your sex partners. A mother with syphilis can give it to her baby during pregnancy or have a miscarriage. Can cause heart disease, brain damage, blindness and death. 	YES
<i>Trichomoniasis ("Trich") or NGU</i>	<ul style="list-style-type: none"> Affects both women and men. In men, trich can cause NGU. Symptoms show up 3-14 days after having sex. Many people have no symptoms. <p>Women:</p> <ul style="list-style-type: none"> Itching, burning or irritation in the vagina. Yellow, greenish or gray discharge from the vagina. <p>Men:</p> <ul style="list-style-type: none"> Watery, white drip from the penis. Burning or pain when you urinate (pee). Need to urinate (pee) more often. 	<ul style="list-style-type: none"> Spread during vaginal sex. 	<ul style="list-style-type: none"> You can give trich to your sex partners. Uncomfortable symptoms will continue. Men can get infections in the prostate gland. 	YES